

**Fires, Floods, and Other Extreme Events — How Watershed Processes
Under Climate Change Will Shape Our Coastlines**

Jonathan A. Warrick,^{1,} Amy E. East,¹ Helen Dow¹*

¹ *U.S. Geological Survey Santa Cruz, California 95060*

**corresponding author (jwarrick@usgs.gov)*

INVITED REVIEW ARTICLE

Cambridge Prisms: Coastal Futures

Journal Resubmit Copy

25rd June 2022

This draft manuscript is distributed solely for purposes of scientific peer review. Its content is deliberative and predecisional, so it must not be disclosed or released by reviewers. Because the manuscript has not yet been approved for publication by the U.S. Geological Survey (USGS), it does not represent any official USGS finding or policy.

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

10.1017/cft.2022.1

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Impact Statement

The future of the world's coasts, including the natural resources and human settlements, will be influenced by ongoing sea-level rise and forthcoming storms, which will combine to increase the likelihood for coastal flooding and erosion. However, many coastal settings receive natural supplies of sediment from adjacent rivers and landscapes, and these sediments can physically build up coastal landforms and reduce the potential for erosion and flooding. It is very important to better understand sediment delivery to the coast, as this is one of the key factors in assessing climate change impacts on coasts. The delivery of new sediment to the coast varies greatly with time, with long intervals with very little sediment discharged punctuated by brief events with tremendous sediment discharge. These sediment input events are generally caused by disturbances in the landscape, which can include wildfires, earthquakes, heavy rainfall, volcanic eruptions, and human impacts. Over decades to centuries, the infrequent, high-magnitude sediment input events can be responsible for most of the sediment that exists along the coast. Climate change has the potential to increase sediment discharge to some coastal settings because of its imminent effects on wildfire, heavy rainfall, and the extent of frozen areas within high altitudes and in polar regions. Because of this, there is increased need to understand the amount of sediment input to coastal regions both now and in the future. Teams of scientists will be needed to monitor and predict future sediment inputs with an eye on better understanding how the coast will respond to climate change.

Abstract

Ongoing sea level rise has brought renewed focus on terrestrial sediment supply to the coast because of its strong influence on whether and how long beaches, marshes, and other coastal landforms may persist into the future. Here we summarize findings of sediment discharge from several coastal rivers, revealing that infrequent, large-magnitude events have disproportionate influence on the morphodynamics of coastal landforms and littoral cells. These event-dominated effects are most pronounced for small, steep mountainous rivers that supply beach and wetland sediment along the world's active tectonic margins, although infrequent events are important drivers of sediment discharge for rivers worldwide. Additionally, extreme events (recurrence intervals of decades to centuries) that follow wildfires, earthquakes, volcanic eruptions, extreme precipitation, or – most notably – combinations of these factors, can redefine coastal sediment budgets and morphology. Some of these extreme events (e.g., wildfires *plus* rainfall) are increasing in magnitude and frequency under modern climate warming, with the likely result of increasing sediment flux to affected coastlines. Climate change is also altering watershed processes in both high latitudes and high altitudes, resulting in increased sediment supply to downstream catchments. We conclude that sediment inputs to coastal systems are highly variable with time, and that the variability and trends in sediment input are as important to characterize as long-term averages.

Keywords

coastal sediment, sea-level rise, accretion, erosion, coastal change, episodic events,
coastal morphodynamics, climate change

Introduction

Coastal landforms and the human communities and natural resources located on these features are increasingly vulnerable to flooding and erosion because of ongoing and accelerating eustatic sea-level rise (FitzGerald et al., 2008; Syvitski et al., 2009; Vitousek et al., 2017; Schuerch et al., 2018). Coastal landforms evolve over time in response to a combination of oceanic, geologic, biologic and fluvial processes, and the relative importance of these processes can vary considerably along the world's coasts and over time (Inman and Nordstrom, 1971; Wright and Short, 1984; Komar, 1998; Friedrichs and Perry, 2001; Murray et al., 2008; Winterwerp et al., 2013; Splinter et al., 2017; Wright et al., 2019). Primary factors in the evolution of many coastal landforms are the volume of sediment available to these systems and the physical processes that move sediment over time (Syvitski, 2005; Frihy et al., 2008; Anthony et al., 2014; Nienhuis et al., 2016; Yang et al., 2017; Tessler et al., 2018; Besset et al., 2019; Warrick et al., 2019).

Coastal landforms can be characterized by the amounts and types of sediment available to them, and there are two endmembers with respect to terrestrial sediment supply: (i) coastal landforms that receive no direct terrestrial sediment, such as many atolls and barrier islands (Heron et al., 1984; FitzGerald et al., 2008; Duvat, 2019), and (ii) landforms that are derived almost entirely from inputs of terrestrial sediment, such as deltas of the world's rivers and littoral cells along many of the world's active

continental margins (Inman and Nordstrom, 1971; Hicks and Inman, 1987; Ashton and Giosan, 2011; Nienhuis et al., 2013; Giosan et al., 2014; Anthony, 2015). For coastal systems that are derived from terrestrial sediment, changes in the volume or grain size of sediment delivered to the coast can result in pronounced changes in the morphology or morphodynamic trajectory (Cooper, 2001; Nienhuis et al., 2013, 2020; Anthony et al., 2014; Giosan et al., 2014; Bendixen et al., 2017; Luo et al., 2017; Warrick et al., 2019; Hoitink et al., 2020; Yang et al., 2020; Syvitski et al., 2022). Thus, to evaluate and predict coastal changes for systems influenced by terrestrial sediment, it is essential to understand several characteristics of the sediment supply, including the processes that deliver sediment to the coast, the timing, volume and characteristics, including grain size distributions, of these sediment contributions, and the littoral processes that disperse sediment from the river mouth (Hicks and Inman, 1987; Orton and Reading, 1993; Kao and Milliman, 2008; Romans et al., 2009; Milliman and Farnsworth, 2013; Nienhuis et al., 2013; Anthony, 2015; Warrick, 2020).

The goal of this paper is to examine and summarize the temporal variability in terrestrial sediment supply to the coast over the decadal to century time scales important to coastal land management. We focus on littoral systems that are derived from terrestrial sources of sediment, which are generally found along active continental margins of the world and have small, steep coastal watersheds that are collectively the dominant supply of sediment to the world's coasts (Inman and Nordstrom, 1971; Milliman and Syvitski, 1992; Milliman and Farnsworth, 2013). Because of the efficient transfer of sediment through these small, steep watersheds (Milliman and Farnsworth, 2013; Romans et al., 2016), we will highlight how coastal supplies of sediment are influenced by natural hazards, infrequent events, and ongoing and pending climate

change. We acknowledge that the sediment budgets of many coastal systems, such as barrier islands and atolls, are not influenced by these watershed processes, owing to negligible sediment contributions from terrestrial landscapes to these coastal landforms (Meade, 1982; Woodroffe et al., 2007; Perry et al., 2015). We also acknowledge that terrestrial supplies of sediment may not be fully integrated into littoral cells, because of offshore transport to the deep sea either during the river discharge event or in the years, decades or centuries following these events (Mulder et al., 2003; Khripounoff et al., 2009; Casalbore et al., 2011; Liu et al., 2016; Romans et al., 2016; Steel et al., 2016; Warrick, 2020). As such, we are reminded that every river and coastal setting is unique, and that care should be taken to understand sediment sources, dispersal processes and time scales, and sinks for each coastal system. In the synthesis below, we summarize characteristic shoreline changes caused by time-varying river sediment discharge and provide considerations for future research and monitoring in the light of ongoing climate change and sea-level rise.

River Sediment Discharge to the Coast

There are abundant examples of fluvial sediment discharge events influencing coastal sediment budgets and morphodynamics. One of these examples is observed at the Rio Rimac of Peru (**Fig. 1**), which discharges exceptional amounts of sediment to the coast during years of high precipitation (French and Mechler, 2017; Guzman et al., 2020). During these wet years, the shoreline at the river mouth progrades 100s of meters seaward, resulting in a river mouth delta with several distributary channels (**Fig. 1a-d**). Following progradation, the delta recedes because of decreases in river sediment discharge and northward littoral sediment transport from wave action (**Fig. 1e-j**). The

littoral transport causes beach widening at least 7 km downdrift of the Rio Rimac mouth (Guzman et al., 2020). Similar patterns of coastal progradation followed by recession and sediment spreading are observed at river mouths around the world as they respond to time-varying sediment inputs (Inman and Nordstrom, 1971; Hicks and Inman, 1987; Cooper, 1993; Anthony and Blivi, 1999; Barnard and Warrick, 2010; Casalbore et al., 2011; Giosan et al., 2012; Milliman and Farnsworth, 2013; Anthony et al., 2014; Bendixen et al., 2017; Besset et al., 2017; Luo et al., 2017; East et al., 2018; Warrick et al., 2019). Satellite imagery can provide important observations of the spatial and temporal response of shorelines to new contributions of river sediment (Besset et al., 2016, 2019; Guzman et al., 2020; Warrick et al., 2022), and example satellite records of the multitude of coastal systems with active river sediment supplies are provided in **Fig. 2**.

Unusually large sediment discharge events can produce shoreline changes that persist for decades or centuries, as evidenced by coastal accretion and geomorphic changes caused by upland volcanic activity on sediment transport from the Santo Tomas River of the Philippines and the Rio Salamá of Guatemala (**Fig. 3**). Coastal accretion at the mouths of both systems extended 100s of meters to several kilometers offshore of the pre-eruption shorelines, and sediment inputs influenced shoreline positions for several to tens of kilometers along the coast (Kuenzi et al., 1979; Siringan and Ringor, 2007). The shoreline accretion from the eruption of Santa Maria, Guatemala continues to extend more than 1 km offshore of the pre-eruption shoreline even though the volcanic event occurred over 120 years ago, providing evidence for the longevity of coastal impacts from rare, but massive, sediment discharge events (Kuenzi et al., 1979; **Fig. 3b**).

Detailed accounting of shoreline responses to river inputs can be obtained where physical monitoring or remote sensing records are adequately frequent (Hicks and Inman, 1987; Barnard and Warrick, 2010; Besset et al., 2016, 2019; Vos et al., 2019a; Guzman et al., 2020; Warrick et al., 2022). For example, the combination of satellite-derived shoreline positions from Landsat and Sentinel-2 imagery using the CoastSat technique (Vos et al., 2019b) and estimates of littoral-grade sediment discharge (Barnard and Warrick, 2010) for the Santa Clara River, California shows how the shoreline responds differently in space and time to elevated river sediment discharge (**Fig. 4**). Sand discharge by the Santa Clara River is punctuated by several wet years, including 1983, 1993, 1995, 1998 and 2005, which combined contributed almost 20 Mt, or approximately 13 million m³, of littoral-grade sand to the coast (**Fig. 4a,b**). The shoreline at the Santa Clara River mouth accreted rapidly during these wet years, followed by multiple year to decadal-scale shoreline recession toward previous positions (**Fig. 4c,d**). In contrast, shorelines greater than 1000 m downcoast from the river mouth had progressively lagged and muted accretion responses (**Fig. 4e-g**). Overall, the beach 2000 m downcoast of the river mouth accreted approximately 60 m between 1990 and 2020 as a result of sediment spreading from the combined river sediment discharge events of the 1990s and 2005 (**Fig. 4e,f**) (Barnard and Warrick, 2010). Combined, these river mouth systems show that sediment input signals can vary greatly in time and that these signals may propagate across and along the shoreline, as described in more detail by coastal observations and theory (Komar, 1973; Hicks and Inman, 1987; Inman et al., 2005; Casalbore et al., 2011; Anthony, 2015; East et al., 2018; Besset et al., 2019; Warrick, 2020).

Temporal Variability in River Sediment Discharge

Infrequent river sediment discharge events that dramatically alter coastal morphology and shoreline positions – such as the floods on the Rio Rímac (**Fig. 1**) and the Santa Clara River (**Fig. 4**), volcanic-related sediment discharge shown for the Santo Tomas River and Rio Salamá (**Fig. 3**), or profound accretion from storm-induced debris-flow activity, as Casalbore et al. (2011) documented on the coast of Sicily – provide important examples of how temporal variations in terrestrial sediment supply can strongly influence littoral systems over spatial scales of kilometers to tens of kilometers and temporal scales of years to over a century. As such, we will explore a few examples of the temporal variability of river sediment discharge, and especially the role of infrequent large discharge events, in the export of sediment from the land to the sea.

To assist with this exploration, we have included a series of sediment discharge records from watersheds ranging from a small, steep river draining the rugged Big Sur coast of California to the world's largest river, the Amazon (**Fig. 5**). Sediment discharge records from these rivers reveal that the smaller watersheds are generally more punctuated by infrequent high discharge events, whereas the massive Amazon River has relatively constant sediment discharge from year to year. Additionally, the records also highlight the effects of perturbations, such as wildfires, floods, earthquakes, typhoons, and human impacts, on year-to-year variations in sediment discharge (**Fig. 5**), which are described more fully in original investigations of these and other rivers (Dadson et al., 2004; Gran and Montgomery, 2005; Wang et al., 2007; Hovius et al., 2011; Lee et al., 2015; Gray, 2018; Montanher et al., 2018; Collins et al., 2020; Warrick et al., 2022). Sediment discharge is especially elevated when a landscape perturbing event, such as wildfire or an earthquake, is followed by heavy precipitation, such as shown in the

records from the Big Sur River and Choshoi River watersheds (**Fig. 5a,b**) (Dadson et al., 2004; Hovius et al., 2011; Warrick et al., 2022). Over larger watershed scales of 100,000s to millions of square kilometers, events such as earthquakes or wildfires impact relatively small areas compared to the river's total drainage basin area and thus contribute only marginally to the overall temporal variations in sediment fluxes (Milliman and Farnsworth, 2013; Francis et al., 2022). Thus, temporal variability in sediment discharge from the large rivers of the world is often attributed to factors that influence broader areas of these watersheds, such as widespread land use change from agriculture development, dams on the mainstem river, or climate patterns influencing the hydrology of the broader basin (Walling, 2006; Wang et al., 2007; Zheng et al., 2018; Syvitski et al., 2022) (**Fig. 5c**).

For the four rivers used in our example, the cumulative sediment discharge of the smaller rivers is more heavily dictated by infrequent events than the larger rivers. This is shown by the steepness of the cumulative sediment discharge curves in **Fig. 5e**, which provides contrast between small rivers such as the Big Sur River of California, for which sediment discharge during the two biggest years represented roughly two-thirds the 50-yr sediment discharge to the coast, and the Amazon River, for which the sediment discharged every year is relatively constant. The high temporal variability in the Big Sur River is largely related to the combined effects of two wildfire and heavy precipitation events (labeled “Fire + Flood”; **Fig. 5a**) (Warrick et al., 2022), although sediment discharge from this river is still strongly variable in time if these events are not considered in the records (**Fig. 5e**). The inverse relationship between watershed-size and temporal variability in river sediment discharge shown in **Figure 5** is consistent with broader understanding of the erosion and sediment transport for rivers throughout

the world (Hicks et al., 2000; Dadson et al., 2004; Kao and Milliman, 2008; Gonzalez-Hidalgo et al., 2010; Milliman and Farnsworth, 2013; Gray, 2018). As such, small, steep watersheds may be considered “reactive” with respect to perturbing effects on sediment discharge, whereas large, continental-scale watersheds may be considered more “buffered” against these effects (*cf.* Romans et al., 2016)

The accounting of sediment discharge to the coast – such as shown in Figure 5 – is generally derived from sampling of river sediment fluxes and applications of models derived from these data (Walling and Fang, 2003; Milliman and Farnsworth, 2013; Syvitski et al., 2022). Unfortunately, the length of river sampling records are generally limited to years to several decades (Milliman and Farnsworth, 2013; Warrick and Milliman, 2018). Although monitoring records are essential for identifying rates and trends in river sediment transport (Gray, 2018), the largest historical events may not be captured by limited duration of sediment sampling. In fact, the exclusion of the largest sediment discharge events is a primary factor for why river sampling records may underestimate long-term watershed sediment yields (Kirchner et al., 2001; Covault et al., 2013). Combined, this suggests that long-term sediment discharge to the coast – especially from the globally important small, steep rivers – is primarily related to the magnitude and frequency of rare large events.

Climate Change

Climate change is modifying the event frequency and intensity of several watershed sediment yield factors discussed above, including the amount and intensity of precipitation and the size, frequency and intensity of wildfires (Westerling et al., 2006; Pachauri et al., 2015; Sankey et al., 2017; Swain et al., 2018; Ball et al., 2021; Touma et

al., 2022). Additionally, rising temperatures are changing the hydrology and sediment yields of both Arctic and alpine landscapes (Bendixen et al., 2017; Li et al., 2021a; Irrgang et al., 2022; Vergara et al., 2022). As such, there is growing understanding that climate change is causing in fundamental changes to the rate of sediment delivery from many landscapes to fluvial and coastal landforms. These effects are most clearly evident in Arctic settings, where terrestrial sediment inputs have increased at the same time that ice-free conditions in the adjacent seas are getting longer (Bendixen et al., 2017; Irrgang et al., 2022). Combined, this is resulting in the expansion of some Arctic deltas – such as those along the Greenland coast – and accelerated erosion on many wave-exposed Arctic shorelines (Bendixen et al., 2017; Irrgang et al., 2022). There is also growing evidence for changes in fluvial sediment discharge in lower latitudes, as wildfire and precipitation are actively changing with climate (Lee et al., 2015; East and Sankey, 2020; Touma et al., 2022). Climate-induced changes are expected to continue with time, and they may dramatically alter sediment budgets of rivers and their downstream coasts.

Conceptual Model and Future Directions

As highlighted above, infrequent fluvial sediment discharge events are a driving factor for many coastal littoral systems worldwide. This time-dependent variability commonly results in wet conditions delivering considerably more sediment than dry conditions do, and when these wet conditions are combined with increases in hillslope sediment supplies from wildfires, earthquakes or volcanic activity, sediment discharge can be exceptional. That is, infrequent events are generally responsible for the majority of sediment transport to the coast (Milliman and Farnsworth, 2013).

We have integrated these concepts into a conceptual model of hypothetical watershed sediment yields and shoreline positions for a small, steep watershed that efficiently conveys sediment from source regions to the coast (*cf.* Roman et al., 2016) that will be used to discuss historic and future trajectories of coastlines (**Fig. 6**). In this simple model there are several perturbations to watershed sediment yield: precipitation, which causes landscape erosion and downstream flooding; wildfire, which denudes the landscape and increases the potential for soil erosion; earthquakes, which liberate regolith throughout the watershed; and volcanic activity, which introduces new sediment materials and disrupts the watershed landscape (**Fig. 6**). As highlighted above, the effects of wildfires on watershed sediment yield are enhanced when they are followed closely by heavy precipitation, as denoted by three “*fire+flood*” events in the hypothetical records (**Fig. 6**). Additionally, the conceptual model includes characteristic human impacts to sediment yields, including increases from land use such as agriculture and road building and decreases from the construction of dams (Kosmas et al., 1997; Vörösmarty et al., 2003; Syvitski, 2005; Walling, 2006; Anthony et al., 2014; Luo et al., 2017; Li et al., 2021b; Syvitski et al., 2022).

The time-varying rate of sediment yield, driven in large part by infrequent events and human impacts, has direct effects on the shoreline position of the littoral cell near the river mouth (**Fig. 6**). For our hypothetical system, small increases in sediment yield result in short-lived changes in the shoreline position, much like the coastal response to monitored events in the Santa Clara River (*cf.* **Fig. 4**). The five large sediment yield events (highlighted with gray shading; **Fig. 6**) cause fundamental changes in the shoreline position as shown by accretion events (arrows; **Fig. 6**). These large events are conceptually similar to the volcanic events highlighted above (*cf.* **Fig. 3**). Lastly, human

impacts may result in coastal accretion or erosion trends, depending on the nature and scale of these impacts (**Fig. 6**).

A few additional items are emphasized in the conceptual model. Firstly, although the river sampling records (blue shading; **Fig. 6**) are shown to capture several decades of sediment discharge including an era of human impacts, they do not include some of the largest and most significant events of the past two centuries. This is rather common for actual river records given that they are commonly years to decades in length (Warrick and Milliman, 2018). To properly understand century-scale or longer sediment yields, river sampling records should be integrated with either an understanding of the role, magnitude, and frequency of events that are not included in the sampling record, or with broader geologic understanding of the discharge record from measurements such as sediment cores or cosmogenic nuclides (Walling, 1988; Lamoureux, 2000; Kirchner et al., 2001; von Blanckenburg, 2005; Covault et al., 2013). Additionally, the lack of exceptional events in most sampling records highlights the importance of records that do include these unique events (e.g., Fan et al., 2019; Gran and Montgomery, 2005; Korup, 2012; Kuenzi et al., 1979; Ritchie et al., 2018; Warrick et al., 2022), largely because they allow for the new understanding to be transferred to longer time scales and to other river systems. Secondly, in looking toward a future with continued climate change (yellow shading; **Fig. 6**), the sustainability of coastal landforms such as the littoral systems highlighted here will depend on whether sediment contributions to the coast can balance the erosion caused by sea-level rise and associated impacts (FitzGerald et al., 2008; Syvitski et al., 2009; Giosan et al., 2014; Nienhuis et al., 2018; Reimann et al., 2018; Schuerch et al., 2018; Hoitink et al., 2020). That is, the future of the world's shorelines will be determined not only by global eustatic sea-level changes,

but also many local factors such as the rate and variability of sediment supply and the conditions and processes that are responsible for transporting sediment throughout the coastal zone (Casalbore et al., 2011; Anthony, 2015; Nienhuis et al., 2016; Steel et al., 2016; Caldwell et al., 2019; Warrick, 2020).

Summarizing, the future of many coasts will be tied to terrestrial sediment inputs (Anthony and Blivi, 1999; Syvitski et al., 2009; Giosan et al., 2012, 2014; Anthony et al., 2014), so a better understanding of the magnitude, frequency, and implications of sediment supply rates is needed. This is especially true for infrequent events, which as noted above can dominate long-term coastal sediment budgets and can redefine coastal morphology. As coastal communities have multiple management options to confront the challenges of climate change, including actively nourishing landforms with imported sediment (de Schipper et al., 2016; Ludka et al., 2018; Armstrong and Lazarus, 2019), these will need to be balanced with an understanding of the inherent coastal processes and morphodynamics, including sediment supplies. In some cases, watershed sediment supplies may be adequate to produce relatively constant shoreline positions for decades or more; in other cases, sediment discharge may be inadequate to keep up with sea-level rise, and coastal erosion and land loss will ensue (**Fig. 6**).

To build this understanding under the current and future conditions of climate change requires collaborative communication and research efforts across hydrologic, geomorphic, and coastal research groups. We encourage the continued development and progress of cross-disciplinary studies of coastal landforms, especially with respect to the linkages between watershed processes and coastal morphodynamics. Although this requires integration across several discipline boundaries (hydrology, oceanography, geomorphology, ecology, meteorology, climate science), it is essential to build this cross-

disciplinary understanding where terrestrial and coastal systems are integrally linked. Additionally, because this work will be highly relevant for coastal communities worldwide that are actively confronting the effects of climate change through land use and expenditure decisions, it is valuable to integrate stakeholder collaboration and the challenges that coastal managers face into research goals and methods (Lemos et al., 2018; Ulibarri et al., 2020). Although the coming era will provide considerable uncertainty for coastal communities and their natural resources, it is crucial that coastal scientists continue to develop relevant information and understanding about our changing coasts.

Acknowledgements

We are thankful for the support of the U.S. Geological Survey (USGS) of this work, including the USGS Coastal and Marine Hazards and Resources Program and a USGS Mendenhall postdoctoral fellowship to H.D. Peter Swarzenski provided comments on an earlier version of the manuscript.

Author Contribution Statement

All authors provided contributions to the development, organization, and writing of this article. J.A.W. compiled data and imagery and drafted the figures.

Financial Support

This work was supported by the U.S. Geological Survey's Coastal and Marine Hazards and Resources Program and the U.S. Geological Survey's Mendenhall Research Fellowship Program.

378

379 **Conflict of Interest**

380 None

381

382 **Data Availability**

383 The data that support the findings of this study are openly available from scientific
 384 reports and publications (from Kuenzi et al., 1979, Wang et al., 2007, Barnard and
 385 Warrick, 2010, Lee et al., 2015, Montanher et al., 2018, and Warrick et al., 2022), the
 386 imagery database contained within Google Earth Pro at
 387 <https://www.google.com/earth/versions/#earth-pro>, and the CoastSat shoreline
 388 position database at <http://coastsat.wrl.unsw.edu.au/>.

389

390 **REFERENCES**

- 391 Anthony, E. J. (2015). Wave influence in the construction, shaping and destruction of
 392 river deltas: A review. *Mar. Geol.* 361, 53–78. doi: 10.1016/j.margeo.2014.12.004.
- 393 Anthony, E. J., and Blivi, A. B. (1999). Morphosedimentary evolution of a delta-sourced,
 394 drift-aligned sand barrier–lagoon complex, western Bight of Benin. *Mar. Geol.*
 395 158, 161–176. doi: 10.1016/S0025-3227(98)00170-4.
- 396 Anthony, E. J., Marriner, N., and Morhange, C. (2014). Human influence and the
 397 changing geomorphology of Mediterranean deltas and coasts over the last 6000
 398 years: From progradation to destruction phase? *Earth-Sci. Rev.* 139, 336–361.
 399 doi: 10.1016/j.earscirev.2014.10.003.
- 400 Armstrong, S. B., and Lazarus, E. D. (2019). Masked Shoreline Erosion at Large Spatial
 401 Scales as a Collective Effect of Beach Nourishment. *Earths Future* 7, 74–84. doi:
 402 10.1029/2018EF001070.
- 403 Ashton, A. D., and Giosan, L. (2011). Wave-angle control of delta evolution: wave-angle
 404 control of delta evolution. *Geophys. Res. Lett.* 38, L13405. doi:
 405 10.1029/2011GL047630.

- 406 Ball, G., Regier, P., González-Pinzón, R., Reale, J., and Van Horn, D. (2021). Wildfires
407 increasingly impact western US fluvial networks. *Nat. Commun.* 12, 2484. doi:
408 10.1038/s41467-021-22747-3.
- 409 Barnard, P. L., and Warrick, J. A. (2010). Dramatic beach and nearshore morphological
410 changes due to extreme flooding at a wave-dominated river mouth. *Mar. Geol.*
411 271, 131–148. doi: 10.1016/j.margeo.2010.01.018.
- 412 Bendixen, M., Lønsmann Iversen, L., Anker Bjørk, A., Elberling, B., Westergaard-
413 Nielsen, A., Overeem, I., et al. (2017). Delta progradation in Greenland driven by
414 increasing glacial mass loss. *Nature* 550, 101–104. doi: 10.1038/nature23873.
- 415 Besset, M., Anthony, E. J., and Bouchette, F. (2019). Multi-decadal variations in delta
416 shorelines and their relationship to river sediment supply: An assessment and
417 review. *Earth-Sci. Rev.* 193, 199–219. doi: 10.1016/j.earscirev.2019.04.018.
- 418 Besset, M., Anthony, E. J., Brunier, G., and Dussouillez, P. (2016). Shoreline change of
419 the Mekong River delta along the southern part of the South China Sea coast
420 using satellite image analysis (1973–2014). *Géomorphologie Relief Process.*
421 *Environ.* 22, 137–146. doi: 10.4000/geomorphologie.11336.
- 422 Besset, M., Anthony, E. J., and Sabatier, F. (2017). River delta shoreline reworking and
423 erosion in the Mediterranean and Black Seas: the potential roles of fluvial
424 sediment starvation and other factors. *Elem Sci Anth* 5, 54. doi:
425 10.1525/elementa.139.
- 426 Caldwell, R. L., Edmonds, D. A., Baumgardner, S., Paola, C., Roy, S., and Nienhuis, J. H.
427 (2019). A global delta dataset and the environmental variables that predict delta
428 formation on marine coastlines. *Earth Surf. Dyn.* 7, 773–787. doi:
429 <https://doi.org/10.5194/esurf-7-773-2019>.
- 430 Casalbore, D., Chiocci, F. L., Scarascia Mugnozza, G., Tommasi, P., and Sposato, A.
431 (2011). Flash-flood hyperpycnal flows generating shallow-water landslides at
432 Fiumara mouths in Western Messina Strait (Italy). *Mar. Geophys. Res.* 32, 257–
433 271. doi: 10.1007/s11001-011-9128-y.
- 434 Collins, B. D., Oakley, N. S., Perkins, J. P., East, A. E., Corbett, S. C., and Hatchett, B. J.
435 (2020). Linking Mesoscale Meteorology With Extreme Landscape Response:
436 Effects of Narrow Cold Frontal Rainbands (NCFR). *J. Geophys. Res. Earth Surf.*
437 125. doi: 10.1029/2020JF005675.
- 438 Cooper, J. A. G. (1993). Sedimentation in a river dominated estuary. *Sedimentology* 40,
439 979–1017. doi: 10.1111/j.1365-3091.1993.tb01372.x.
- 440 Cooper, J. A. G. (2001). Geomorphological variability among microtidal estuaries from
441 the wave-dominated South African coast. *Geomorphology* 40, 99–122. doi:
442 10.1016/S0169-555X(01)00039-3.

- 443 Covault, J. A., Craddock, W. H., Romans, B. W., Fildani, A., and Gosai, M. (2013).
 444 Spatial and Temporal Variations in Landscape Evolution: Historic and Longer-
 445 Term Sediment Flux through Global Catchments. *J. Geol.* 121, 35–56. doi:
 446 10.1086/668680.
- 447 Dadson, S. J., Hovius, N., Chen, H., Dade, W. B., Lin, J.-C., Hsu, M.-L., et al. (2004).
 448 Earthquake-triggered increase in sediment delivery from an active mountain belt.
 449 *Geology* 32, 733. doi: 10.1130/G20639.1.
- 450 de Schipper, M. A., de Vries, S., Ruessink, G., de Zeeuw, R. C., Rutten, J., van Gelder-
 451 Maas, C., et al. (2016). Initial spreading of a mega feeder nourishment:
 452 Observations of the Sand Engine pilot project. *Coast. Eng.* 111, 23–38. doi:
 453 10.1016/j.coastaleng.2015.10.011.
- 454 Duvat, V. K. E. (2019). A global assessment of atoll island planform changes over the
 455 past decades. *WIREs Clim. Change* 10. doi: 10.1002/wcc.557.
- 456 East, A. E., and Sankey, J. B. (2020). Geomorphic and Sedimentary Effects of Modern
 457 Climate Change: Current and Anticipated Future Conditions in the Western
 458 United States. *Rev. Geophys.* 58. doi: 10.1029/2019RG000692.
- 459 East, A. E., Stevens, A. W., Ritchie, A. C., Barnard, P. L., Campbell-Swarzenski, P.,
 460 Collins, B. D., et al. (2018). A regime shift in sediment export from a coastal
 461 watershed during a record wet winter, California: Implications for landscape
 462 response to hydroclimatic extremes. *Earth Surf. Process. Landf.* 43, 2562–2577.
 463 doi: 10.1002/esp.4415.
- 464 Fan, X., Scaringi, G., Korup, O., West, A. J., Westen, C. J., Tanyas, H., et al. (2019).
 465 Earthquake-Induced Chains of Geologic Hazards: Patterns, Mechanisms, and
 466 Impacts. *Rev. Geophys.* 57, 421–503. doi: 10.1029/2018RG000626.
- 467 FitzGerald, D. M., Fenster, M. S., Argow, B. A., and Buynevich, I. V. (2008). Coastal
 468 Impacts Due to Sea-Level Rise. *Annu. Rev. Earth Planet. Sci.* 36, 601–647. doi:
 469 10.1146/annurev.earth.35.031306.140139.
- 470 Francis, O., Fan, X., Hales, T., Hobley, D., Xu, Q., and Huang, R. (2022). The Fate of
 471 Sediment After a Large Earthquake. *J. Geophys. Res. Earth Surf.* 127. doi:
 472 10.1029/2021JF006352.
- 473 French, A., and Mechler, R. (2017). Managing El Niño Risks Under Uncertainty in Peru:
 474 Learning from the past for a more disaster-resilient future. Available at:
 475 [http://pure.iiasa.ac.at/id/eprint/14849/1/French_Mechler_2017_El%20Ni%C3](http://pure.iiasa.ac.at/id/eprint/14849/1/French_Mechler_2017_El%20Ni%C3%20B1o_Risk_Peru_Report.pdf)
 476 [%B1o_Risk_Peru_Report.pdf](http://pure.iiasa.ac.at/id/eprint/14849/1/French_Mechler_2017_El%20Ni%C3%20B1o_Risk_Peru_Report.pdf) [Accessed January 31, 2022].
- 477 Friedrichs, C. T., and Perry, J. E. (2001). Tidal salt marsh morphodynamics: a synthesis.
 478 *J. Coast. Res.*, 7–37.

- 479 Frihy, O. E., Shereet, S. M., and El Banna, M. M. (2008). Pattern of Beach Erosion and
 480 Scour Depth along the Rosetta Promontory and their Effect on the Existing
 481 Protection Works, Nile Delta, Egypt. *J. Coast. Res.* 244, 857–866. doi:
 482 10.2112/07-0855.1.
- 483 Giosan, L., Coolen, M. J. L., Kaplan, J. O., Constantinescu, S., Filip, F., Filipova-
 484 Marinova, M., et al. (2012). Early Anthropogenic Transformation of the Danube-
 485 Black Sea System. *Sci. Rep.* 2. doi: 10.1038/srep00582.
- 486 Giosan, L., Syvitski, J. P. M., Constantinescu, S., and Day, John (2014). Protecting the
 487 world's deltas. *Nature* 516, 31–33.
- 488 Gonzalez-Hidalgo, J. C., Batalla, R. J., Cerdà, A., and de Luis, M. (2010). Contribution of
 489 the largest events to suspended sediment transport across the USA. *Land*
 490 *Degrad. Dev.* 21, 83–91. doi: 10.1002/ldr.897.
- 491 Gran, K. B., and Montgomery, D. R. (2005). Spatial and temporal patterns in fluvial
 492 recovery following volcanic eruptions: Channel response to basin-wide sediment
 493 loading at Mount Pinatubo, Philippines. *GSA Bull.* 117, 195–211. doi:
 494 10.1130/B25528.1.
- 495 Gray, A. B. (2018). The impact of persistent dynamics on suspended sediment load
 496 estimation. *Geomorphology* 322, 132–147. doi:
 497 10.1016/j.geomorph.2018.09.001.
- 498 Guzman, E., Ramos, C., and Dastgheib, A. (2020). Influence of the El Niño Phenomenon
 499 on Shoreline Evolution. Case Study: Callao Bay, Perú. *J. Mar. Sci. Eng.* 8, 90. doi:
 500 10.3390/jmse8020090.
- 501 Heron, S. D., Moslow, T. F., Berelson, W. M., Herbert, J. R., Steele, G. A., and Susman,
 502 K. R. (1984). "Holocene Sedimentation of A Wave-Dominated Barrier-Island
 503 Shoreline: Cape Lookout, North Carolina," in *Developments in Sedimentology*
 504 *Hydrodynamics and Sedimentation in Wave-Dominated Coastal Environments.*,
 505 eds. B. Greenwood and R. A. Davis (Elsevier), 413–434. doi: 10.1016/S0070-
 506 4571(08)70157-2.
- 507 Hicks, D. M., Gomez, B., and Trustrum, N. A. (2000). Erosion thresholds and
 508 suspended sediment yields, Waipaoa River Basin, New Zealand. *Water Resour.*
 509 *Res.* 36, 1129–1142. doi: 10.1029/1999WR900340.
- 510 Hicks, D. M., and Inman, D. L. (1987). Sand dispersion from an ephemeral river delta on
 511 the Central California coast. *Mar. Geol.* 77, 305–318. doi: 10.1016/0025-
 512 3227(87)90119-8.
- 513 Hoitink, A. J. F., Nittrouer, J. A., Passalacqua, P., Shaw, J. B., Langendoen, E. J.,
 514 Huismans, Y., et al. (2020). Resilience of River Deltas in the Anthropocene. *J.*
 515 *Geophys. Res. Earth Surf.* 125. doi: 10.1029/2019JF005201.

- 516 Hovius, N., Meunier, P., Lin, C.-W., Chen, H., Chen, Y.-G., Dadson, S., et al. (2011).
 517 Prolonged seismically induced erosion and the mass balance of a large
 518 earthquake. *Earth Planet. Sci. Lett.* 304, 347–355. doi:
 519 10.1016/j.epsl.2011.02.005.
- 520 Inman, D. L., Jenkins, S. A., McLachlan, A., Orme, A. R., Leatherman, S. P., Whitman,
 521 D., et al. (2005). “Accretion and Erosion Waves on Beaches,” in *Encyclopedia of*
 522 *Coastal Science*, ed. M. L. Schwartz (Dordrecht: Springer Netherlands), 1–116.
 523 doi: 10.1007/1-4020-3880-1_1.
- 524 Inman, D. L., and Nordstrom, C. E. (1971). On the tectonic and morphologic
 525 classification of coasts. *J. Geol.* 79, 1–21.
- 526 Irrgang, A. M., Bendixen, M., Farquharson, L. M., Baranskaya, A. V., Erikson, L. H.,
 527 Gibbs, A. E., et al. (2022). Drivers, dynamics and impacts of changing Arctic
 528 coasts. *Nat. Rev. Earth Environ.* 3, 39–54. doi: 10.1038/s43017-021-00232-1.
- 529 Kao, S. J., and Milliman, J. D. (2008). Water and Sediment Discharge from Small
 530 Mountainous Rivers, Taiwan: The Roles of Lithology, Episodic Events, and
 531 Human Activities. *J. Geol.* 116, 431–448. doi: 10.1086/590921.
- 532 Khripounoff, A., Vangriesheim, A., Crassous, P., and Etoubleau, J. (2009). High
 533 frequency of sediment gravity flow events in the Var submarine canyon
 534 (Mediterranean Sea). *Mar. Geol.* 263, 1–6. doi: 10.1016/j.margeo.2009.03.014.
- 535 Kirchner, J. W., Finkel, R. C., Riebe, C. S., Granger, D. E., Clayton, J. L., King, J. G., et
 536 al. (2001). Mountain erosion over 10 yr, 10 ky, and 10 my time scales. *Geology*
 537 29, 591–594.
- 538 Komar, P. D. (1973). Computer Models of Delta Growth due to Sediment Input from
 539 Rivers and Longshore Transport. *Geol. Soc. Am. Bull.* 84, 2217–2226.
- 540 Komar, P. D. (1998). *Beach processes and sedimentation*. 2nd ed. New York: Prentice
 541 Hall.
- 542 Korup, O. (2012). Earth’s portfolio of extreme sediment transport events. *Earth-Sci.*
 543 *Rev.* 112, 115–125. doi: 10.1016/j.earscirev.2012.02.006.
- 544 Kosmas, C., Danalatos, N., Cammeraat, L. H., Chabart, M., Diamantopoulos, J., Farand,
 545 R., et al. (1997). The effect of land use on runoff and soil erosion rates under
 546 Mediterranean conditions. *CATENA* 29, 45–59. doi: 10.1016/S0341-
 547 8162(96)00062-8.
- 548 Kuenzi, W. D., Horst, O. H., and McGehee, R. V. (1979). Effect of volcanic activity on
 549 fluvial-deltaic sedimentation in a modern arc-trench gap, southwestern
 550 Guatemala. *Geol. Soc. Am. Bull.* 90, 827–838.

- 551 Lamoureux, S. (2000). Five centuries of interannual sediment yield and rainfall-induced
 552 erosion in the Canadian High Arctic recorded in lacustrine varves. *Water Resour.*
 553 *Res.* 36, 309–318. doi: 10.1029/1999WR900271.
- 554 Lee, T.-Y., Huang, J.-C., Lee, J.-Y., Jien, S.-H., Zehetner, F., and Kao, S.-J. (2015).
 555 Magnified Sediment Export of Small Mountainous Rivers in Taiwan: Chain
 556 Reactions from Increased Rainfall Intensity under Global Warming. *PLOS ONE*
 557 10, e0138283. doi: 10.1371/journal.pone.0138283.
- 558 Lemos, M. C., Arnott, J. C., Ardoin, N. M., Baja, K., Bednarek, A. T., Dewulf, A., et al.
 559 (2018). To co-produce or not to co-produce. *Nat. Sustain.* 1, 722–724. doi:
 560 10.1038/s41893-018-0191-0.
- 561 Li, D., Lu, X., Overeem, I., Walling, D. E., Syvitski, J., Kettner, A. J., et al. (2021a).
 562 Exceptional increases in fluvial sediment fluxes in a warmer and wetter High
 563 Mountain Asia. *Science* 374, 599–603. doi: 10.1126/science.abi9649.
- 564 Li, M., Li, T., Zhu, L., Meadows, M. E., Zhu, W., and Zhang, S. (2021b). Effect of Land
 565 Use Change on Gully Erosion Density in the Black Soil Region of Northeast China
 566 From 1965 to 2015: A Case Study of the Kedong County. *Front. Environ. Sci.* 9,
 567 652933. doi: 10.3389/fenvs.2021.652933.
- 568 Liu, J. T., Hsu, R. T., Hung, J.-J., Chang, Y.-P., Wang, Y.-H., Rendle-Bühring, R. H., et
 569 al. (2016). From the highest to the deepest: The Gaoping River–Gaoping
 570 Submarine Canyon dispersal system. *Earth-Sci. Rev.* 153, 274–300. doi:
 571 10.1016/j.earscirev.2015.10.012.
- 572 Ludka, B. C., Guza, R. T., and O'Reilly, W. C. (2018). Nourishment evolution and
 573 impacts at four southern California beaches: A sand volume analysis. *Coast. Eng.*
 574 136, 96–105. doi: 10.1016/j.coastaleng.2018.02.003.
- 575 Luo, X. X., Yang, S. L., Wang, R. S., Zhang, C. Y., and Li, P. (2017). New evidence of
 576 Yangtze delta recession after closing of the Three Gorges Dam. *Sci. Rep.* 7. doi:
 577 10.1038/srep41735.
- 578 Meade, R. H. (1982). Sources, Sinks, and Storage of River Sediment in the Atlantic
 579 Drainage of the United States. *J. Geol.* 90, 235–252.
- 580 Milliman, J. D., and Farnsworth, K. L. (2013). *River discharge to the coastal ocean: a*
 581 *global synthesis*. Cambridge University Press.
- 582 Milliman, J. D., and Syvitski, J. P. M. (1992). Geomorphic/tectonic control of sediment
 583 discharge to the ocean: The importance of small mountainous rivers. *J. Geol.* 100,
 584 525–544. doi: 10.1086/629606.
- 585 Montanher, O. C., Novo, E. M. L. de M., and Souza Filho, E. E. de (2018). Temporal
 586 trend of the suspended sediment transport of the Amazon River (1984–2016).
 587 *Hydrol. Sci. J.* 63, 1901–1912. doi: 10.1080/02626667.2018.1546387.

- 588 Mulder, T., Syvitski, J. P. M., Migeon, S., Faugères, J.-C., and Savoye, B. (2003). Marine
589 hyperpycnal flows: initiation, behavior and related deposits. A review. *Mar. Pet.*
590 *Geol.* 20, 861–882. doi: 10.1016/j.marpetgeo.2003.01.003.
- 591 Murray, A. B., Knaapen, M. A. F., Tal, M., and Kirwan, M. L. (2008).
592 Biomorphodynamics: Physical-biological feedbacks that shape landscapes:
593 OPINION. *Water Resour. Res.* 44. doi: 10.1029/2007WR006410.
- 594 Nienhuis, J. H., Ashton, A. D., Edmonds, D. A., Hoitink, A. J. F., Kettner, A. J.,
595 Rowland, J. C., et al. (2020). Global-scale human impact on delta morphology
596 has led to net land area gain. *Nature* 577, 514–518. doi: 10.1038/s41586-019-
597 1905-9.
- 598 Nienhuis, J. H., Ashton, A. D., Nardin, W., Fagherazzi, S., and Giosan, L. (2016).
599 Alongshore sediment bypassing as a control on river mouth morphodynamics. *J.*
600 *Geophys. Res. Earth Surf.* 121, 664–683. doi: 10.1002/2015JF003780.
- 601 Nienhuis, J. H., Ashton, A. D., Roos, P. C., Hulscher, S. J., and Giosan, L. (2013). Wave
602 reworking of abandoned deltas. *Geophys. Res. Lett.* 40, 5899–5903.
- 603 Nienhuis, J. H., Hoitink, A. J. F. (Ton), and Törnqvist, T. E. (2018). Future Change to
604 Tide-Influenced Deltas. *Geophys. Res. Lett.* 45, 3499–3507. doi:
605 10.1029/2018GL077638.
- 606 Orton, G. J., and Reading, H. G. (1993). Variability of deltaic processes in terms of
607 sediment supply, with particular emphasis on grain size. *Sedimentology* 40, 475–
608 512. doi: 10.1111/j.1365-3091.1993.tb01347.x.
- 609 Pachauri, R. K., Mayer, L., and Intergovernmental Panel on Climate Change eds. (2015).
610 *Climate change 2014: synthesis report*. Geneva, Switzerland: Intergovernmental
611 Panel on Climate Change.
- 612 Perry, C. T., Kench, P. S., O’Leary, M. J., Morgan, K. M., and Januchowski-Hartley, F.
613 (2015). Linking reef ecology to island building: Parrotfish identified as major
614 producers of island-building sediment in the Maldives. *Geology* 43, 503–506.
615 doi: 10.1130/G36623.1.
- 616 Reimann, L., Vafeidis, A. T., Brown, S., Hinkel, J., and Tol, R. S. J. (2018).
617 Mediterranean UNESCO World Heritage at risk from coastal flooding and
618 erosion due to sea-level rise. *Nat. Commun.* 9, 4161. doi: 10.1038/s41467-018-
619 06645-9.
- 620 Ritchie, A. C., Warrick, J. A., East, A. E., Magirl, C. S., Stevens, A. W., Bountry, J. A., et
621 al. (2018). Morphodynamic evolution following sediment release from the world’s
622 largest dam removal. *Sci. Rep.* 8. doi: 10.1038/s41598-018-30817-8.

- 623 Romans, B. W., Castelltort, S., Covault, J. A., Fildani, A., and Walsh, J. P. (2016).
 624 Environmental signal propagation in sedimentary systems across timescales.
 625 *Earth-Sci. Rev.* 153, 7–29. doi: 10.1016/j.earscirev.2015.07.012.
- 626 Romans, B. W., Normark, W. R., McGann, M. M., Covault, J. A., and Graham, S. A.
 627 (2009). Coarse-grained sediment delivery and distribution in the Holocene Santa
 628 Monica Basin, California: Implications for evaluating source-to-sink flux at
 629 millennial time scales. *Geol. Soc. Am. Bull.* 121, 1394–1408. doi:
 630 10.1130/B26393.1.
- 631 Sankey, J. B., Kreitler, J., Hawbaker, T. J., McVay, J. L., Miller, M. E., Mueller, E. R., et
 632 al. (2017). Climate, wildfire, and erosion ensemble foretells more sediment in
 633 western USA watersheds: Future Fire and Sediment. *Geophys. Res. Lett.* 44,
 634 8884–8892. doi: 10.1002/2017GL073979.
- 635 Schuerch, M., Spencer, T., Temmerman, S., Kirwan, M. L., Wolff, C., Lincke, D., et al.
 636 (2018). Future response of global coastal wetlands to sea-level rise. *Nature* 561,
 637 231–234. doi: 10.1038/s41586-018-0476-5.
- 638 Siringan, F. P., and Ringor, C. L. (2007). Changes in the Position of the Zambales
 639 Shoreline Before and After the 1991 Mt. Pinatubo Eruption: Controls of Shoreline
 640 Change. in.
- 641 Splinter, K. D., Turner, I. L., Reinhardt, M., and Ruessink, G. (2017). Rapid adjustment
 642 of shoreline behavior to changing seasonality of storms: observations and
 643 modelling at an open-coast beach: Adjustment of shoreline behavior to changing
 644 seasonality of storms. *Earth Surf. Process. Landf.* 42, 1186–1194. doi:
 645 10.1002/esp.4088.
- 646 Steel, E., Simms, A. R., Warrick, J., and Yokoyama, Y. (2016). Highstand shelf fans: The
 647 role of buoyancy reversal in the deposition of a new type of shelf sand body. *Geol.*
 648 *Soc. Am. Bull.* 128, 1717–1724. doi: 10.1130/B31438.1.
- 649 Swain, D. L., Langenbrunner, B., Neelin, J. D., and Hall, A. (2018). Increasing
 650 precipitation volatility in twenty-first-century California. *Nat. Clim. Change* 8,
 651 427–433. doi: 10.1038/s41558-018-0140-y.
- 652 Syvitski, J., Ángel, J. R., Saito, Y., Overeem, I., Vörösmarty, C. J., Wang, H., et al.
 653 (2022). Earth's sediment cycle during the Anthropocene. *Nat. Rev. Earth*
 654 *Environ.* doi: 10.1038/s43017-021-00253-w.
- 655 Syvitski, J. P. M. (2005). Impact of humans on the flux of terrestrial sediment to the
 656 global coastal ocean. *Science* 308, 376–380. doi: 10.1126/science.1109454.
- 657 Syvitski, J. P. M., Kettner, A. J., Overeem, I., Hutton, E. W. H., Hannon, M. T.,
 658 Brakenridge, G. R., et al. (2009). Sinking deltas due to human activities. *Nat.*
 659 *Geosci.* 2, 681–686. doi: 10.1038/ngeo629.

- 660 Tessler, Z. D., Vörösmarty, C. J., Overeem, I., and Syvitski, J. P. M. (2018). A model of
 661 water and sediment balance as determinants of relative sea level rise in
 662 contemporary and future deltas. *Geomorphology* 305, 209–220. doi:
 663 10.1016/j.geomorph.2017.09.040.
- 664 Touma, D., Stevenson, S., Swain, D. L., Singh, D., Kalashnikov, D. A., and Huang, X.
 665 (2022). Climate change increases risk of extreme rainfall following wildfire in the
 666 western United States. *Sci. Adv.* 8, eabmo320. doi: 10.1126/sciadv.abmo320.
- 667 Ulibarri, N., Goodrich, K. A., Wagle, P., Brand, M., Matthew, R., Stein, E. D., et al.
 668 (2020). Barriers and opportunities for beneficial reuse of sediment to support
 669 coastal resilience. *Ocean Coast. Manag.* 195, 105287. doi:
 670 10.1016/j.ocecoaman.2020.105287.
- 671 Vergara, I., Garreaud, R., and Ayala, Á. (2022). Sharp Increase of Extreme Turbidity
 672 Events Due To Deglaciation in the Subtropical Andes. *J. Geophys. Res. Earth*
 673 *Surf.* 127. doi: 10.1029/2021JF006584.
- 674 Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L., and Storlazzi, C. D.
 675 (2017). Doubling of coastal flooding frequency within decades due to sea-level
 676 rise. *Sci. Rep.* 7, 1399.
- 677 von Blanckenburg, F. (2005). The control mechanisms of erosion and weathering at
 678 basin scale from cosmogenic nuclides in river sediment. *Earth Planet. Sci. Lett.*
 679 237, 462–479. doi: 10.1016/j.epsl.2005.06.030.
- 680 Vörösmarty, C. J., Meybeck, M., Fekete, B., Sharma, K., Green, P., and Syvitski, J. P. M.
 681 (2003). Anthropogenic sediment retention: major global impact from registered
 682 river impoundments. *Glob. Planet. Change* 39, 169–190. doi: 10.1016/S0921-
 683 8181(03)00023-7.
- 684 Vos, K., Harley, M. D., Splinter, K. D., Simmons, J. A., and Turner, I. L. (2019a). Sub-
 685 annual to multi-decadal shoreline variability from publicly available satellite
 686 imagery. *Coast. Eng.* 150, 160–174. doi: 10.1016/j.coastaleng.2019.04.004.
- 687 Vos, K., Splinter, K. D., Harley, M. D., Simmons, J. A., and Turner, I. L. (2019b).
 688 CoastSat: A Google Earth Engine-enabled Python toolkit to extract shorelines
 689 from publicly available satellite imagery. *Environ. Model. Softw.* 122, 104528.
 690 doi: 10.1016/j.envsoft.2019.104528.
- 691 Walling, D. E. (1988). Erosion and sediment yield research — Some recent perspectives.
 692 *J. Hydrol.* 100, 113–141. doi: 10.1016/0022-1694(88)90183-7.
- 693 Walling, D. E. (2006). Human impact on land–ocean sediment transfer by the world’s
 694 rivers. *Geomorphology* 79, 192–216. doi: 10.1016/j.geomorph.2006.06.019.

- 695 Walling, D. E., and Fang, D. (2003). Recent trends in the suspended sediment loads of
 696 the world's rivers. *Glob. Planet. Change* 39, 111–126. doi: 10.1016/S0921-
 697 8181(03)00020-1.
- 698 Wang, H., Yang, Z., Saito, Y., Liu, J. P., Sun, X., and Wang, Y. (2007). Stepwise
 699 decreases of the Huanghe (Yellow River) sediment load (1950–2005): Impacts of
 700 climate change and human activities. *Glob. Planet. Change* 57, 331–354. doi:
 701 10.1016/j.gloplacha.2007.01.003.
- 702 Warrick, J. A. (2020). Littoral Sediment From Rivers: Patterns, Rates and Processes of
 703 River Mouth Morphodynamics. *Front. Earth Sci.* 8, 355. doi:
 704 10.3389/feart.2020.00355.
- 705 Warrick, J. A., and Milliman, J. D. (2018). Do we know how much fluvial sediment
 706 reaches the sea? Decreased river monitoring of U.S. coastal rivers. *Hydrol.*
 707 *Process.* 32, 3561–3567. doi: 10.1002/hyp.13276.
- 708 Warrick, J. A., Stevens, A. W., Miller, I. M., Harrison, S. R., Ritchie, A. C., and
 709 Gelfenbaum, G. (2019). World's largest dam removal reverses coastal erosion.
 710 *Sci. Rep.* 9, 1–12. doi: 10.1038/s41598-019-50387-7.
- 711 Warrick, J. A., Vos, K., East, A. E., and Vitousek, S. (2022). Fire (plus) flood (equals)
 712 beach: coastal response to an exceptional river sediment discharge event. *Sci.*
 713 *Rep.* 12, 3848. doi: 10.1038/s41598-022-07209-0.
- 714 Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W. (2006). Warming
 715 and Earlier Spring Increase Western U.S. Forest Wildfire Activity. *Science* 313,
 716 940–943. doi: 10.1126/science.1128834.
- 717 Winterwerp, J. C., Erftemeijer, P. L. A., Suryadiputra, N., van Eijk, P., and Zhang, L.
 718 (2013). Defining Eco-Morphodynamic Requirements for Rehabilitating Eroding
 719 Mangrove-Mud Coasts. *Wetlands* 33, 515–526. doi: 10.1007/s13157-013-0409-x.
- 720 Woodroffe, C. D., Samosorn, B., Hua, Q., and Hart, D. E. (2007). Incremental accretion
 721 of a sandy reef island over the past 3000 years indicated by component-specific
 722 radiocarbon dating. *Geophys. Res. Lett.* 34, L03602. doi:
 723 10.1029/2006GL028875.
- 724 Wright, L. D., and Short, A. D. (1984). Morphodynamic variability of surf zones and
 725 beaches: A synthesis. *Mar. Geol.* 56, 93–118. doi: 10.1016/0025-3227(84)90008-
 726 2.
- 727 Wright, L. D., Syvitski, J. P. M., Nichols, C. R., and Zinnert, J. (2019). “Coastal
 728 Morphodynamics and Ecosystem Dynamics,” in *Tomorrow's Coasts: Complex*
 729 *and Impermanent Coastal Research Library.*, eds. L. D. Wright and C. R. Nichols
 730 (Cham: Springer International Publishing), 69–84. doi: 10.1007/978-3-319-
 731 75453-6_5.

- 732 Yang, H. F., Yang, S. L., Xu, K. H., Wu, H., Shi, B. W., Zhu, Q., et al. (2017). Erosion
733 potential of the Yangtze Delta under sediment starvation and climate change. *Sci.*
734 *Rep.* 7. doi: 10.1038/s41598-017-10958-y.
- 735 Yang, S. L., Luo, X., Temmerman, S., Kirwan, M., Bouma, T., Xu, K., et al. (2020). Role
736 of delta-front erosion in sustaining salt marshes under sea-level rise and fluvial
737 sediment decline. *Limnol. Oceanogr.* 65, 1990–2009. doi: 10.1002/lno.11432.
- 738 Zheng, S., Xu, Y. J., Cheng, H., Wang, B., Xu, W., and Wu, S. (2018). Riverbed erosion of
739 the final 565 kilometers of the Yangtze River (Changjiang) following construction
740 of the Three Gorges Dam. *Sci. Rep.* 8. doi: 10.1038/s41598-018-30441-6.

741

Figure Captions

Figure 1. The influence of river sediment discharge on the coastal morphology and shoreline positions at the mouth of Rio Rímac, Peru during 2016-2021. As described by Guzman et al. (2020), heavy flooding in early 2017 resulted in massive growth of the river mouth delta and spreading of this sediment northward in the subsequent years, similar to the coastal morphodynamics following flooding in 1983 and 1998. Imagery from Google Earth.



Figure 2. Examples of coastal changes at the mouths of small rivers of the world resulting from contributions of new sediment. Imagery from Google Earth.



Figure 3. Decadal to century persistence of coastal accretion from increases in river sediment yield resulting from volcanic activity in coastal watersheds. (a) The mouth of the Santo Tomas River 28 yr after the eruption of Mount Pinatubo, Philippines. (b) The mouth of Rio Salamá almost 120 yr after the eruption of Santa Maria, Guatemala. Additional shorelines from before and immediately following the eruptions from publicly available Landsat imagery or interpretations of Kuenzi et al. (1979). Imagery from Google Earth.



Figure 4. River sediment discharge and shoreline positions of the Santa Clara River, California highlighting the effects of infrequent events on shoreline accretion and the spatial and temporal variations of shoreline response to new sediment. (a) Annual rainfall at a National Weather Service station near the river. (b) Littoral-grade sand (>125 micron) discharge from the Santa Clara River after Barnard and Warrick (2010); data from 2009-2021 were not estimated due to a lack of river gaging. (c-g) Shoreline positions from five transects derived from CoastSat analyses of Vos et al. (2019b). Shoreline positions are normalized to the average position of each transect during 1990-1992 when the shoreline was consistently narrow. (h) Satellite imagery of the Santa Clara River mouth following the 2005 sediment discharge events from Google Earth. Locations of the shoreline from a September 2004 image and the CoastSat transects are shown.




Figure 5. Annual sediment discharge measurements for four different rivers highlighting how temporal variations are influenced by perturbations such as wildfires, floods and earthquakes, and the size of the watershed. Time series shown in (a) to (d) have been transformed into ranked annual exceedance values in (e) using the cumulative sediment discharge measured in each river. Recurrence intervals were estimated by the reciprocal of the annual exceedance probabilities. Data for (a) to (d) were derived from Warrick et al. (2022), Lee et al. (2015), Wang et al. (2007), and Montanher et al. (2018), respectively. Descriptive terms about the watershed sizes (righthand side) are derived from discussion in Romans et al. (2016).




Figure 6. Conceptual model of coastal responses to watershed processes for a theoretical small, steep river basin (after the fire-flood model of Keller et al., 1997). Relative sediment yield of the watershed is influenced by stochastic events, including floods, wildfires, earthquakes, volcanic activity, and combined events, such as wildfire followed by flooding (“F+F”). The shoreline position of the littoral cell responds to increases in watershed sediment yield with accretion events (upward pointing arrows), because of the efficient transfer of river sediment to the littoral cell. Future shoreline positions (righthand side) will be determined by balance between sediment supply and sea-level rise. Also highlighted are hypothetical intervals of river sampling, climate change effects, land-use-change effects, and damming of the river.

